P44 42 MAPK(ERK1 2) Mouse Monoclonal Antibody(6B1) Catalog No.: RA10050

Basic Information

	nforma				4 ·		
lnt		PP	n	വ	П	U.	n
, , , , , ,	w			a i	"	U.	ш

Reactivity H,M,R

Immunogen Synthetic Peptide

Host Mouse

Isotype IgG1

Storage Buffer & Condition 1mg/ml in PBS, pH 7.4, containing 0.02% sodium

azide and 50% glycerol.

Observed MW 44,42KD

Applications Recommended Dilution

WB 1:1,000-2,000

IHC 1:100-200


Preparation & Storage


Storage Storage Storage Storage

shipment.

Shipping Bule Ice

Experimental Data





Western blot analysis of 1)Mouse Brain Immunohistochemical Tissue, 2)Rat Brain Tissue with P44/42 paraffin-embedded Hu MAPK(ERK1/2)Mouse mAb diluted at using P44/42 MAPK 1:2,000. diluted at 1:200.

Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma using P44/42 MAPK (ERK1/2)Mouse mAb diluted at 1:200.

Background

Mitogen-activated protein kinases (MAPKs) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (Erk1/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli including mitogens, growth factors, and cytokines and is an important target in the diagnosis and treatment of cancer.