HO-1 Mouse Monoclonal Anbtibody(3F9)

Catalog No.: RA10301

Basic Information

		4 •
\mathbf{I}	nrm	ation
		auun

Reactivity H,M,R

Immunogen Recombinant Protein

Host Mouse

Isotype IgG

Storage Buffer & Condition 1mg/ml in PBS, pH 7.4, containing 0.02% sodium

azide and 50% glycerol.

Observed MW ~28KD

Applications Recommended Dilution

WB 1:1,000-2,000


Preparation & Storage

Storage Storage Storage Storage

shipment.

Shipping Bule Ice

Experimental Data



Western blot analysis of 3T3 Cell Lysate using HO-1 Mouse Monoclonal mAb diluted at 1:2,000.

Background

Catalytic activity:Heme + 3 AH(2) + 3 O(2) = biliverdin + Fe(2+) + CO + 3 A + 3 H(2)O.,function:Heme oxygenase cleaves the heme ring at the alpha methene bridge to form biliverdin. Biliverdin is subsequently converted to bilirubin by biliverdin reductase. Under physiological conditions,the activity of heme oxygenase is highest in the spleen,where senescent erythrocytes are sequestrated and destroyed.,induction:Heme oxygenase 1 activity is highly inducible by its substrate heme and by various non-heme substances such as heavy metals,bromobenzene,endotoxin,oxidizing agents and UVA.,similarity:Belongs to the heme oxygenase family.