# Cleaved PARP Mouse Monoclonal Antibody(M4) Catalog No.: RA10398

#### **Basic Information**

т	P		4 ·
$\mathbf{I}$	tor	mai	tion
	ш	ша	UUU

**Reactivity** H

**Immunogen** Synthetic Peptide

**Host** Mouse

Isotype IgG1

Storage Buffer & Condition 1mg/ml in PBS, pH 7.4, containing 0.02% sodium

azide and 50% glycerol.

Observed MW 116,89KD

A 180 / 0	T 1 1 T 1 1 T 1 1 1 1 1 1 1 1 1 1 1 1 1
<b>Applications</b>	<b>Recommended Dilution</b>

**WB** 1:1,000-3,000

IHC 1:200-500


## **Preparation & Storage**


Storage Storage Storage Storage

shipment.

**Shipping** Bule Ice

## **Experimental Data**





Western blot analysis of 1)Jurkat ,2)Hela, with Cleaved PARP Mouse mAb diluted at 1:2,000.

Immunohistochemical analysis of paraffin-embedded Human Tonsil Tissue using Cleaved PARP Mouse mAb diluted at 1:500.

#### **Background**

Poly [ADP-ribose] polymerase 1 (PARP-1) also known as NAD+ ADP-ribosyltransferase 1 or poly[ADP-ribose] synthase 1 is an enzyme that in humans is encoded by the PARP1 gene. PARP1 has a role in repair of single-stranded DNA (ssDNA) breaks. Knocking down intracellular PARP1 levels with siRNA or inhibiting PARP1 activity with small molecules reduces repair of ssDNA breaks. In the absence of PARP1, when these breaks are encountered during DNA replication, the replication fork stalls, and double-strand DNA (dsDNA) breaks accumulate.