MAPK(ERK1 2) Rabbit Polyclonal Antibody(F136) Catalog No.: RA20277

Basic Information

т	r	forma			
In.		rn	10	f I	on
	w		1a	u	UII

Reactivity H,M,R

Immunogen Recombinant Protein

Host Rabbit

Isotype IgG

Storage Buffer & Condition 1mg/ml in PBS, pH 7.4, containing 0.02% sodium

azide and 50% glycerol.

Observed MW 44,42KD

WB 1:1,000-2,000

IHC 1:50-100


Preparation & Storage


Storage Storage Storage Storage

shipment.

Shipping Bule Ice

Experimental Data





Western blot analysis of 1)HepG2 Cell Lysate, 2)3T3 Cell Lysate using ERK1/2 Rabbit pAb diluted at 1:2000

Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma Tissue using ERK1/2 Rabbit pAb diluted at 1:100

Background

Mitogen-activated protein kinases (MAPKs) are a widely conserved family of serine/threonine protein kinases involved in many cellular programs such as cell proliferation, differentiation, motility, and death. The p44/42 MAPK (Erk1/2) signaling pathway can be activated in response to a diverse range of extracellular stimuli including mitogens, growth factors, and cytokines and is an important target in the diagnosis and treatment of cancer.