# **Recombinant Human/Mouse/Rat BMP-2**


# Catalog No.: RP0040

## **Basic Information**

| Information               |   |
|---------------------------|---|
| Source                    | E.coli  |
| Description               | Recombinant Human/Mouse/Rat Bone Morphogenetic Protein 2 is produced by our E.coli expression system and the target gene encoding Gln283-Arg396 is expressed.   |
| Accession                 | P12643  |
| Known As                  | Bone Morphogenetic Protein 2; BMP-2; Bone Morphogenetic Protein 2A; BMP-2A; BMP2; BMP2A   |
| <b>Predicted Mol Mass</b> | 13.3 KDa  |
| <b>Apparent Mol Mass</b>  | 13 KDa, reducing conditions   |
| Properties                |   |
| Formulation               | Lyophilized from a 0.2 µm filtered solution of 10mM HAc-NH <sub>4</sub> Ac, 4% D-Mannitol, pH 4.0.  |
| Storage                   | Lyophilized protein should be stored at $\leq -20$ °C, stable for one year after receipt.<br>Reconstituted protein solution can be stored at 2-8 °C for 2-7 days.<br>Aliquots of reconstituted samples are stable at $\leq -20$ °C for 3 months.  |
| Endotoxin                 | $< 0.01 \text{ EU}/\mu g$ as determined by LAL test.  |
| Reconstitution            | Always centrifuge tubes before opening.Do not mix by vortex or pipetting.<br>It is not recommended to reconstitute to a concentration less than 100µg/ml.<br>Dissolve the lyophilized protein in 50mM Acetic Acid.<br>Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |
| Shipping                  | The product is shipped at ambient temperature.<br>Upon receipt, store it immediately at the temperature listed below.   |

#### **Reed Biotech Ltd**

### **Experimental Data**



# Purity-SDS-PAGE

Greater than 95% as determined by reducing SDS-PAGE. (QC verified)

## Background

Bone Morphogenetic Protein-2 (BMP-2) is one of the bone-growth regulatory factors that belong to the transforming growth factor-beta (TGF-beta) superfamily of proteins. BMPs are synthesized as large precursor molecules, which are cleaved by proteolytic enzymes. The active form of BMP-2 can consist of a dimer of two identical proteins or a heterodimer of two related bone morphogenetic proteins.