

## Recombinant Mouse FLT3LG (C-6His)

Catalog No.: RP0090

### Basic Information

#### Information


<b>Source</b>	<i>Human Cells</i>
<b>Description</b>	Recombinant Mouse Fms-like Tyrosine Kinase 3 Ligand is produced by our Mammalian expression system and the target gene encoding Gly27-Arg188 is expressed with a 6His tag at the C-terminus.
<b>Accession</b>	P49772
<b>Known As</b>	Fms-related tyrosine kinase 3 ligand (Flt3L); SL cytokine; Flt3 ligand;
<b>Predicted Mol Mass</b>	19.4 KDa
<b>Apparent Mol Mass</b>	22-30 KDa, reducing conditions

#### Properties


<b>Formulation</b>	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
<b>Storage</b>	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$ , stable for one year after receipt. Reconstituted protein solution can be stored at $2-8^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
<b>Endotoxin</b>	$< 1 \text{ EU}/\mu\text{g}$ as determined by LAL test.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\mu\text{g}/\text{ml}$ . Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.

## Experimental Data

### Purity-SDS-PAGE



### Bioactivity-BLI



Greater than 95% as determined by reducing SDS-PAGE. (QC verified)

Loaded Recombinant Human FLT3 (C-Fc) on Pro-A Biosensor, can bind Recombinant Mouse FLT3LG (C-6His) with an affinity constant of 0.447 nM as determined in BLI assay. (Regularly tested)

## Background

Fms-related tyrosine kinase 3 ligand(Flt3L) is a single-pass type I membrane protein and consists of 232 amino acids. Flt3L is a hematopoietic four helical bundle cytokine, structurally homologous to stem cell factor and colony stimulating factor. Flt3L synergizes well with a number of other colony stimulating factors and interleukins. Flt3L stimulates the proliferation and differentiation of various blood cell progenitors by activating FLT3.